

The brief but long history of X-Ray Polarimetry

Martin C. Weisskopf
(NASA/MSFC)

Tuesday May 24, 2016

Outline

- A look to the past
- The present
- A brief view to the future

In the beginning.....

- July 1968 – Lithium-block, Thomson-scattering polarimeter flown on an Aerobee -150 rocket
 - Target was Sco X-1






Fig. 1. (a) Schematic representation of the polarimeter concept. (b) Mounting of the polarimeter and ancillary equipment in the rocket.

Scattering polarimeter


- Thomson cross-section illustrates the angular dependence

$$d\sigma / d\Omega = (e^2 / mc^2)^2 (\cos^2 \theta \cos^2 \varphi + \sin^2 \varphi)$$


- For scattering from bound electrons one must account for both coherent and incoherent scattering and photoelectric absorption



Thompson Approximation



Cos (polar scattering angle)



Azimuthal scattering angle

Advantages

- Inherently broad band device
 - Astrophysical non-thermal spectra are characteristically broad band
 - Astrophysical diagnostics (model discriminators) should benefit from understanding the energy dependence

Considerations

- Scatter as much incident flux as possible
- Avoid multiple scattering
- Collect as many scattered X-Rays as possible
- Minimize the background
- Achieve as large a sensitivity to polarization as possible
 - Optimize the “MDP” at the 99% confidence level

$$MDP(\%) = (4.29 \times 10^4 / M(\%)) \sqrt{(R_S + R_B)} / \sqrt{R_S^2 t}$$

Disadvantage: The Polarimeter Conundrum


- The scattering material should be thick (deep) in order to effectively provide for interaction with all the incident photons.
- The scattering material should be thin (narrow) in order to allow the scattered photon to easily escape.
- Similar conundrums apply as well to other approaches to X-ray polarimetry.

In the beginning....


- March 1969 - Lithium-block, Thomson-scattering polarimeter flown on an Aerobee - 150 rocket
 - Target was the Crab Nebula
- February 1971 Lithium-block, Thomson-scattering polarimeter and a Bragg crystal polarimeter flown on an Aerobee -350 rocket
 - Target was the Crab Nebula
- Three rockets in 21 months!

Rocket 17.09

- Two instruments in one payload!
 - Lithium scattering polarimeter
 - 4 Bragg crystal polarimeters



Crystal polarimeter





Rocket 17.09

- 1971 Aerobee 350
 - Crab detection!
 - $P = 15\% \pm 5\%$
 - $\phi = 156 \pm 10^\circ$




On to the satellite experiment

- 1975 OSO-8 crystal polarimeter
 - Precision measurement of integrated Crab Nebula polarization at 2.6 keV
 - $P = 19\% \pm 1\%$
 - $\phi = 156^\circ \pm 2^\circ$ (NNE)



Compare to modern detailed optical results

Moran, P., Shearer, A., Mignani, R.P., et al. (2013)



1975 - 2014

- X-ray polarimeter on the original Einstein mission but removed via descoping
 - Polarimeters amongst the first to go
- X-ray polarimeter built for the original SRG mission ... which never launched
- Polarimeters don't get much observatory time
 - 60 days of observation per year (OSO-8)
 - 11 days of observations per year (SXG)
- X-ray polarimeter selected for the last NASA SMEX mission ... but was cancelled


Electron-Tracking Polarimeters

- Optical Imaging Chamber
 - Austin & Ramsey 1992
- Pixelated Gas Multiplication
 - Costa et al. 2001
- Time Projection Chamber
 - Black et al. 2007

- The direction of the *initial* K-shell photoelectron is determined by the electric vector and the direction of the incoming photon


$$\frac{d\sigma}{d\Omega} = f(\zeta) r_0^2 Z^5 \alpha_0^4 \left(\frac{1}{\beta} \right)^{7/2} 4\sqrt{2} \sin^2 \theta \cos^2 \varphi$$

Electron-Tracking



Site of initial
ionization and
Auger electron
cloud produced by
a 54 keV photon

Electron-Tracking



Looking to the future

- We need to remember that X-ray polarimetry is difficult
 - One does not expect all astrophysical systems to be strongly polarized
 - Instruments typically not fully sensitive to polarization
 - Linear polarization is non-negative – i.e. one always measures something, even for an unpolarized source

Looking to the future

- X-ray polarimetry can be powerful – *especially* if accompanied by high-resolution imaging
- X-ray polarimetry typically *requires* longer observing times than imaging, spectroscopy, and timing
- **Do not rely on an observatory-class mission**
 - Won't get the needed observing time
 - Early (if not first) candidate for descoping

- Polarimetry is the study of systematic effects
 - End-to end calibration of the full system with unpolarized beams to the appropriate precision is essential!

A look to the future

- So many missions being planned
 - IXPE (NASA SMEX?) 2020
 - Praxys (NASA SMEX? 2020
 - XIPE (ESA M4?) 2025
 - CHINA
 - JAPAN
 - INDIA